

## Impact of the zebra mussel invasion on the ecological integrity of Lough Sheelin, Ireland: distribution, population characteristics and water quality changes in the lake<sup>\*</sup>

Michael Millane<sup>1\*</sup>, Mary Kelly-Quinn<sup>1</sup> and Trevor Champ<sup>2</sup>

<sup>1</sup>Freshwater Biodiversity, Ecology and Fisheries Research Group, School of Biology and Environmental Science, University College Dublin, Belfield, Dublin 4, Ireland

<sup>2</sup>Central Fisheries Board, Swords Business Campus, Swords, Co. Dublin, Ireland

\*Corresponding author

E-mail: [michael.millane@ucd.ie](mailto:michael.millane@ucd.ie)

Received 26 June 2008; accepted in revised form 29 July 2008; published online 15 August 2008

### Abstract

The zebra mussel, *Dreissena polymorpha* (Pallas 1771), invaded Lough Sheelin in the midlands of Ireland in 2001. In order to assess the status of the mussel population in the lake, the distribution, extent of colonisation, abundance, biomass and size-frequency structure of post-settlement stages were studied in 2005 and 2006. In addition, changes to water quality parameters in the lake post-establishment were assessed with reference to the pre-invasion period. Zebra mussels were found throughout the lake at most sites on all main categories of substrates examined (stony and soft substrate; submerged vegetation: Characeae, *Elodea* spp., *Cladophora* spp., *Potamogeton* spp. and *Myriophyllum* spp., and emergent vegetation: *Phragmites australis* and *Schoenoplectus lacustris*). Overall, increases in density and biomass of mussels were recorded from 2005 to 2006. Analysis of physiochemical data show a reduction in chlorophyll *a* with an increase in water transparency, however the total phosphorus concentration remains high. This paper highlights the common misconception that zebra mussel introductions lead to overall improvements in water quality.

*Key words:* *Dreissena polymorpha*, lakes, population, Sheelin, water quality

### Introduction

The zebra mussel, *Dreissena polymorpha* (Pallas 1771), is one of the most notable alien species to invade aquatic ecosystems in recent times

(Karatayev et al. 2002; Minchin et al. 2002a). It is responsible for many significant ecological and economic impacts throughout freshwater and estuarine waterbodies in both Europe and North America (reviewed in Karatayev et al. 1997; Munawar et al. 2005; Pimentel et al. 2005;

<sup>\*</sup> This paper was presented at the special session "Dreissenology" of the 15th International Conference on Aquatic Invasive Species – ICAIS 2007, September 23-27, 2007, Nijmegen

Lovell et al. 2006). Once introduced, mussel populations can expand rapidly to extensively colonise a system, often becoming established as the dominant benthic invertebrate within a few years after initial invasion. Exponential population growth is typically evident within the first few years of colonisation followed by a levelling off (e.g. Burlakova et al. 2006; Lucy et al. 2005; Lucy 2006). This may be preceded by an initial lag phase where the invasion is not detected (Burlakova et al. 2006). In the longer term, population densities can be relatively unstable, fluctuating by varying degrees from year to year (e.g. Stańczykowska and Lewandowski 1993; Burla and Ribi 1998; Hunter and Simons 2004; Strayer and Malcom 2006). Densities in excess of 120,000 m<sup>-2</sup> and a biomass >5kgm<sup>-2</sup> are possible in lakes (Minchin et al. 2002a; Minchin et al. 2005).

In Ireland, the zebra mussel was introduced around 1994 or earlier (Minchin and Moriarty 1998a; Minchin and Moriarty 1998b). Since then it has extensively colonised many hard fresh-water systems (Minchin et al. 2002b; Minchin 2003). It is believed that the mussel was originally introduced into the lower River Shannon region, being inadvertently transported here attached to leisure craft coming from the UK (Minchin and Moriarty 1998b; Pollux et al. 2003; Astanei et al. 2005). The introduction of the zebra mussel to Lough Sheelin most likely occurred in late 2001 (Minchin et al. 2002b; Kerins et al. 2007).

Zebra mussels predominantly live on hard stony substrates but can successfully inhabit soft sediments as well (Marsden 1992; Marsden and Lansky 2000; Coakley et al. 2002). In addition, mussels can extensively colonise submerged vegetation (Sullivan et al. 2002; Burlakova et al. 2006) and attach to hard artificial structures present in the water (Minchin et al. 2002b; Lancioni and Gaino 2006). In some lakes, particularly those without large areas of stony substrate, submerged and semi-submerged vegetation can provide the principal settlement substrate for the zebra mussel (Lewandowski 1982; Ramcharan et al. 1992; Hunter and Simons 2004). Furthermore, they can settle on other fauna, most notably on the shells of unionid mussels (Lucy et al. 2005; Mackie 1993; Maguire et al. 2003). Irish populations are believed to have a typical lifespan of 2 to 3 years or cease growth after that time (Maguire 2002; Lucy et al. 2005).

The magnitude and extent of ecological impacts are primarily related to the abundance and biomass of the resident zebra mussel population, biomass being most relevant, as it corresponds more directly to filter-feeding capacity (Vanderploeg et al. 2002; Burlakova et al. 2006). Being prolific filter-feeders (Reeders et al. 1993; Karatayev et al. 1997), established mussel populations can remove large amounts of seston, including phytoplankton and small zooplankton (Reeders et al. 1993; Wong et al. 2003), from the water column and deposit it on the bottom as faeces and pseudofaeces (Reeders et al. 1993; Vanderploeg et al. 2002). This can result in the suppression of pelagic primary production (Fahnenstiel et al. 1995; Caraco et al. 1997). The associated increase in water transparency allows light penetration to deeper depths which can promote the expansion of submerged macrophytes (Skubinna et al. 1995; Zhu et al. 2006) and the growth of epilithic algae (Lowe and Pillsbury 1995).

The aims of this study were (1) to determine the distribution, extent of colonisation and the population characteristics of the zebra mussel in Lough Sheelin on hard, soft and plant substrates and (2) to evaluate any changes in the principal water quality parameters in the lake post-colonisation. The establishment of baseline invasion data on the zebra mussel will be used to assist with the formation of a long-term monitoring programme to complement existing fish population and physiochemical datasets managed by the Irish Central Fisheries Board.

## Material and Methods

### *Study sites*

Lough Sheelin (53°48'N, 7°20'W) is situated in the River Inny catchment, a sub-catchment of the River Shannon in the midlands of Ireland. It is a highly alkaline lake (90-220 mg/l CaCO<sub>3</sub>, Champ 1998), underlain by a carboniferous limestone geology. By Irish standards it is a fairly large waterbody with a surface area of 18.5 km<sup>2</sup>, however it is relatively shallow having a maximum depth of 14 metres with approximately 75% of lake less than 6 metres deep (John et al. 1982). The lake floor comprises predominantly of soft substrates such as mud, marl and sand. The shoreline is fringed by stony substrate in all but the southern part of the lake, with more


substantive stony areas present in the eastern portion. Generally these do not extend a great distance from the shore. Submerged vegetation, primarily consisting of charophytes (Family Characeae), *Elodea* spp., *Cladophora* spp. and *Potamogeton* spp., are abundant in the shallow southwesterly section of the lake in addition to the western and southern areas nearer the shoreline and in some other sheltered bays. The reed beds, located around the fringes of the lake are of made up of two main species, *Phragmites australis* (Cav. Trin. ex Steud.) and *Schoenoplectus lacustris* (Linn., Pallas).

Lough Sheelin was originally an excellent wild brown trout fishery of good water quality (Champ 1977). However since the early 1970s, the lake has been under considerable environmental pressure from eutrophication, the primary source of this pollution originated from intensive agricultural practices, particularly from pig farming and the associated spreading of the resulting slurry in the catchment area (Champ 1993). A brief period of recovery ensued in the early 1990s due to reductions in phosphorus

inputs to the lake (Champ 1993). More recently, anthropogenic sources such as local municipal wastewater facilities and local industry may additionally be a contributing factor to the high nutrient load of the lake (Kerins et al. 2007). The lake is particularly suited for zebra mussels, having adequate pH, calcium content, food availability and temperature ranges (Ramcharan et al. 1992; Stańczykowska and Lewandowski 1993; Champ 1998).

#### Sampling and sample processing

To establish distribution, extent of colonisation and population characteristics of the resident zebra mussel population, four principal substrate types (stony substrate, soft substrate, submerged vegetation and reed beds) were examined for settled juvenile ( $\geq 1\text{mm}$ ) and adult stages in spring (stony and soft substrates) and autumn (submerged and reed bed vegetation) of 2005 and 2006 (see Figure 1 for a map of sampling sites). Five sites were sampled in the stony substrate (S1-S5) with five replicates taken at



**Figure 1.** Map of Lough Sheelin showing sites sampled for zebra mussels (▲ stony substrate S1-S5; ● soft substrate G1-G18; ○ submerged vegetation P1-P5; and ■ reed bed vegetation R1-R6)

three distance intervals (10m, 15m and 25m) following a transect line from the shore using a quadrat square (size 25cm x 25cm). Scuba diving or snorkelling was employed for this. At site S5,

due to a lack of adequate stony area, five replicates were randomly taken along a transect line running parallel to the shore at a distance of 5m out (adapted from Lucy 2005). On the soft

substrate, eighteen sites (G1-G18) were sampled for zebra mussels using an Eckman grab (size 15cm x 15cm). Six replicates were taken per site (adapted from Marsden 1992). In both the stony and soft substrate sites, a visual estimation of percentage substrate type was used to classify substrate based on the Wentworth scale (Wentworth 1922). In addition, five submerged vegetation (P1-P5) sites were sampled using a double-ended rake thrown out five times from a boat (adapted from Irvine et al. 2001). Retrieved plant material was separated per taxa encountered (Characeae, *Elodea* spp., *Cladophora* spp., *Potamogeton* spp. and *Myriophyllum* spp.). Furthermore, six reed beds (R1-R6) were randomly selected for examination. These were comprised of *Phragmites australis* and/or *Schoenoplectus lacustris*. Ten random mature specimens of each reed species were individually cut at their base from the outer fringe of the reed bed and carefully removed for further examination (Sullivan et al. 2002). The location of all sites were recorded using a hand held GPS (model Garmin Etrex Summit 3.10).

Tissue blotted dry wet weight was used as a measure for zebra mussel biomass and also for the biomass of submerged vegetation. Mussel density (number/m<sup>2</sup>) and biomass (g/m<sup>2</sup>) were determined at the stony and soft substrate sites. Size frequency distributions were generated by randomly selecting circa two hundred mussels (if present) and measuring each to the nearest millimetre along the longest axis of the shell.

Physiochemical data were provided by the Central Fisheries Board, which has a long-term water monitoring programme on the lake. Generally, monthly water samples were taken from a mid-lake station. Physiochemical analyses were carried out using standard methods as described in Champ (1998) and Kerins et al. (2007).

#### *Substrate mapping and determination of total abundance, total biomass and filtering capacity of mussel population*

A map of the hard and soft substrates in the lake was developed using a combination of direct observations (snorkelling, SCUBA diving, shoreline and boat survey work) and through the grab sampling work. This was integrated with a digitised bathymetric map modified from the Ordnance Survey Discovery Series map 34 and from the bathymetric map in Gargan and O'Grady (1992).

To assess total abundance and biomass of mussels in the lake, mean values were determined per depth interval and proportional area of substrate type, and a weighted average was calculated to produce an overall empirical estimate (adapted from Karatayev et al. 1990 as cited in Lucy 2005; and from Burlakova et al. 2006). As plant sampling for zebra mussels was semi-quantitative, it was not possible to include these in overall calculations of total density and biomass. For overall filtering capacity, an arbitrary rate of 44ml of water filtered per gram of wet total mass per hour was assigned (Karatayev et al. 1997; Lucy et al. 2005). The total volume of water in the lake (88x10<sup>6</sup>m<sup>3</sup>, Champ 1998) was subsequently divided by the daily filtering rate to provide an estimate of the total filtering capacity of zebra mussels in Lough Sheelin.

#### *Statistical analysis*

Potential changes in density and biomass from 2005 to 2006 were evaluated per distance interval and per site using Mann-Whitney U tests. Actual densities were compared to densities predicted using the models in Ramcharan et al. (1992) based on pH, calcium ions, PO<sub>4</sub> and NO<sub>3</sub>. Potential changes to physiochemical parameters (chlorophyll *a*, Secchi disc, total organic nitrogen (TON) and total phosphorus (TP)) were assessed by comparing the pre zebra mussel period (1996-2000) with the post-establishment period (2003-2007) using a nested analysis of variance. The relationship between physiochemical variables was also explored before (1991-2000) and after the zebra mussel (2003-2007) establishment using Spearman's Rank Correlations. Assumptions such as normality and homogeneity of variances were tested a priori as appropriate to the statistical test employed. Statistica 8.0 was used for all statistical analyses (StatSoft Inc. 2008).

#### **Results**

Zebra mussels were present at all stony substrate sites (S1-S5) and at all soft substrate sites < 6m depth (G1-G9) sampled in spring 2005. In spring 2006, mussels were found at a number of additional soft substrate sites up to 9m depth (G10, G12 and G17). No mussels were located during either sampling period at the deepest, centrally located soft substrate site (G15) which

was >9m depth. Stony areas were found to contain densities of zebra mussels three to five times greater than soft substrate sites ( $2157 \pm 416 \text{ m}^{-2}$  versus  $703 \pm 196 \text{ m}^{-2}$  in 2005 and  $3016 \pm 434 \text{ m}^{-2}$  versus  $587.7 \pm 154 \text{ m}^{-2}$  in 2006 respecti-

vely; Table 1). Biomass was also substantially higher on this substrate ( $1522 \pm 270 \text{ gm}^{-2}$  versus  $230.7 \pm 59 \text{ gm}^{-2}$  in 2005 and  $2723 \pm 344 \text{ gm}^{-2}$  versus  $207.6 \pm 58 \text{ gm}^{-2}$  in 2006 respectively; Table 1).

**Table 1.** Characteristics of zebra mussels on the principal substrate types

Year	Parameter	Mean/ Range	Stony substrate	Soft substrate	Characeae	<i>Elodea</i> spp.	<i>S. lacustris</i>
2005	Biomass† ( $\text{g/m}^2$ )	Mean	1522 (270)	230.7 (59)	0.43 (0.18)	0.41 (0.22)	1.3 (0.22)
		Range	4.8-9698	0.89-2842	<0.01-1.6	<0.01-3.84	<0.1-6.3
	Density <sup>1</sup> (no.s/ $\text{m}^2$ )	Mean	2157 (416)	703 (196)	3.6 (1.99)	4.2 (3)	5.98 (1.3)
		Range	32-20288	44.4-11144	0.07-22.7	0.01-52	1-41
	Shell length (mm)	Mean	14.7 (2.3)	9.7 (1.2)	7.4 (1)	8.7 (1.4)	10.6 (0.6)
		Range	3-28	1-26	2-23	2-22	1-22
2006	Biomass† ( $\text{g/m}^2$ )	Mean	2723 (344)	207.6 (58)	0.22 (0.12)	0.09 (0.06)	3 (0.9)
		Range	112-9490	2.2-3605	<0.01-1.83	<0.01-0.63	0.4-30
	Density <sup>1</sup> (no.s/ $\text{m}^2$ )	Mean	3016 (434)	587.7 (154)	1.25 (0.71)	0.66 (0.41)	8 (2.58)
		Range	96-13936	44.4-9679.2	<0.01-10.8	<0.01-4.47	1-84
	Shell length (mm)	Mean	16.2 (2.7)	11.5 (0.95)	10.6 (1.1)	10.6 (0.4)	13.2 (1)
		Range	3-31	2-27	1-18	1-20	2-21

Standard error in brackets; <sup>1</sup>for Characeae and *Elodea* spp.=numbers per g wet weight of plant tissue / for *S. lacustris* = numbers per individual plant; †for Characeae and *Elodea* spp.=biomass per g wet weight of plant tissue / for *S. lacustris* = biomass per individual plant; Minimum value in range is minimum non-zero value

An overall increase in biomass and density were evident on the stony substrates from spring 2005 to spring 2006 (Table 1). However, Mann-Whitney U tests revealed no specific significant differences ( $P>0.05$ ) for mussel biomass or density on the stony substrate per distance interval at each site. Densities did decrease for sites S1 25m, S2 10m and S3 10m, although this was not found to be significant (Mann-Whitney U,  $P>0.05$ ). The highest mean biomass and density were found at site S4 in 2005 ( $3150 \pm 780 \text{ gm}^{-2}$  and  $4851 \pm 1453 \text{ m}^{-2}$  respectively) and the lowest at site S1 ( $186 \pm 54 \text{ gm}^{-2}$  and  $485 \pm 147 \text{ m}^{-2}$  respectively). In 2006, site S5 had the highest mean biomass ( $5013 \pm 1393 \text{ gm}^{-2}$ ) and S4 the highest mean density ( $6402 \pm 1230 \text{ m}^{-2}$ ). Again, the lowest mean biomass ( $589 \pm 272 \text{ gm}^{-2}$ ) and density ( $555 \pm 203 \text{ m}^{-2}$ ) were found at site S1. At all stony sites with distance intervals (S1-S4), biomass and density successively increased with greater distance from shore and with depth (range 0.2m-2.3m).


On the soft substrate, biomass typically decreased at most sites. However, only G9 experienced a significant decline (from  $1630 \pm 534 \text{ gm}^{-2}$  to  $2.22 \pm 2.22 \text{ gm}^{-2}$ , Mann-Whitney U,  $P<0.01$ ), whereas G8 was the only site to have a significant increase in biomass (from  $39.2 \pm 20.95 \text{ gm}^{-2}$  to  $1722 \pm 501 \text{ gm}^{-2}$ , Mann-Whitney

U,  $P<0.01$ ). Density was generally stable on the soft substrates with the only changes corresponding to the trends in biomass for G9 (from  $7208 \pm 2256 \text{ m}^{-2}$  to  $14.8 \pm 14.8 \text{ m}^{-2}$ ; Mann-Whitney U,  $P<0.01$ ) and G8 (from  $141 \pm 72 \text{ m}^{-2}$  to  $3693 \pm 979 \text{ m}^{-2}$ , Mann-Whitney U,  $P<0.01$ ). Indeed, site G9 had the highest biomass and density on the soft substrate in spring 2005. Most other sites had a mean mussel biomass  $<300 \text{ gm}^{-2}$  and a density of  $<800 \text{ m}^{-2}$ , except for sites G2 (from  $1175 \pm 591 \text{ gm}^{-2}$  and  $1243 \pm 591 \text{ m}^{-2}$  respectively) and G6 ( $649 \pm 223 \text{ gm}^{-2}$  and  $1680 \pm 630 \text{ m}^{-2}$  respectively). In spring 2006, the highest biomass and density were recorded at site G8 (see above). For other sites, mean biomass was  $<150 \text{ gm}^{-2}$ , except for sites G1 ( $872 \pm 448 \text{ gm}^{-2}$ ), and G6 ( $539 \pm 171 \text{ gm}^{-2}$ ), and mean density typically  $<130 \text{ m}^{-2}$ , except for sites G1 ( $3012 \pm 1525 \text{ m}^{-2}$ ), G2 ( $466 \pm 357 \text{ m}^{-2}$ ), G6 ( $1724 \pm 441 \text{ m}^{-2}$ ) and G7 ( $836 \pm 645 \text{ m}^{-2}$ ).

Zebra mussels were found attached to each of the five main submerged plant taxa in autumn 2005 and autumn 2006. Mussels were present at all submerged sites, except at site P5 in 2006 where no plant material was retrieved. Characeae and *Elodea* spp. represented the principal substrate for attached mussels in both sampling periods. *Cladophora* spp., *Potamogeton* spp. and *Myriophyllum* spp. were relatively uncolonised,

however these taxa were not found at all sites sampled. Mean biomass of mussels per biomass of plant and number of individuals per biomass of plant declined in 2006 for both Characeae (49% and 65% respectively) and *Elodea* spp. (79% and 84% respectively). As regards reed beds, *Schoenoplectus lacustris* was utilised for settlement to a greater extent than *Phragmites australis*, with no mussels found on the latter in 2006 and only 1 specimen attached in 2005 at site R6 (Table 1).


Clear age cohorts could not be identified from site specific or overall length frequency distributions, except for mussels on the soft substrate in spring 2005 (see Figures 2A-3B) when all data were combined. This was particularly influenced



**Figure 2.** % length frequency of *Dreissena polymorpha* on stony substrates (all sites combined) in spring 2005 (n=5945) (A) and in spring 2006 (n=7300) (B)

by site G9, which comprised 57% of the specimens in the overall distribution. Two modes are evident here, representing two possible age

classes of approximately 1-7 mm and 8-19 mm respectively (Figure 3A). No unimodal distribution common to all sites was evident. Mean shell length on the stony substrates ( $14.7 \pm 2.3$  mm in 2005 and  $16.2 \pm 2.7$  mm in 2006) was greater than on the soft substrates ( $9.7 \pm 1.2$  mm in 2005 and  $11.5 \pm 0.95$  mm in 2006) and on the submerged and reed bed vegetation (Table 1).



**Figure 3.** % length frequency of *Dreissena polymorpha* on soft substrates (all sites combined) in spring 2005 (n=1709) (A) and in spring 2006 (n=1348) (B)

Models from Ramcharan et al. (1992) applied to the dataset predicted the resident zebra mussel population in the lake would be in the low category ( $<3000$  mussels per  $m^2$ ), with densities of mussels of  $2441m^{-2}$  in 2005 and  $2483m^{-2}$  in 2006. For these models, pH was entered as 8.19,  $Ca^{2+}$  as  $41mg/l$  and  $PO_4$  as the average of available readings of  $0.0126 mg/l$  in 2005 and  $0.008 mg/l$  in 2006. Due to a scarcity of information for  $NO_3$ , a value of  $0.417mg/l$  was assumed in both years.

Estimates for total abundance, total biomass and corresponding filtering capacity for zebra mussels on stony and soft substrates in both sampling periods are given in Table 2. These suggest an increase in total abundance of approximately 62% from spring 2005 with a similar increase in total biomass of 71%. The increase in total biomass has resulted in a notable reduction in the estimated theoretical time for the zebra mussel to filter the lake (from 22.4 to 13.1 days).

Analysis of the physiochemical data (Table 3) show a significant reduction in chlorophyll *a* (ANOVA,  $F_{1,86}=39.23$ ,  $P<0.0001$ ) and an increase in Secchi disc values (ANOVA,  $F_{1,81}=60.23$ ,  $P<0.0001$ ), from the period before

zebra mussel colonisation to the time since establishment of the mussel. Although a significant increase is evident for TP concentrations (ANOVA,  $F_{1,70}=11.05$ ,  $P<0.01$ ), these data could not be made homogeneous so the result must be treated with caution. Nested ANOVAs are robust to departures from homocedasticity when there are many independent estimates of residual variability (Underwood 1997). However, as only three readings were available for this variable in 2000, this precluded the use of other appropriate statistical tests to evaluate changes to TP. No significant change was detected for TON (ANOVA,  $F_{1,82}=0.96$ ,  $P>0.05$ ).

**Table 2.** Total abundance, biomass and filtration capacity of zebra mussels in Lough Sheelin

Year	Total abundance <sup>1</sup>	Total biomass <sup>1</sup> (kg)	Filtration capacity (m <sup>3</sup> /day)	Estimated time to filter lake (days)
2005	$10.36 \times 10^9$	$3.73 \times 10^6$	$3.93 \times 10^6$	22.4
2006	$16.38 \times 10^9$	$6.36 \times 10^6$	$6.71 \times 10^6$	13.1

<sup>1</sup> stony and soft substrates

**Table 3.** Mean values for physiochemical parameters before and after establishment of the zebra mussel

Parameter	Pre-zebra mussel (1996-2000)	Post-zebra mussel (2003-2007)	Significance
Chlorophyll <i>a</i> (mg/m <sup>3</sup> )	22.5 (1.96)	10.19 (1.69)	$P<0.0001$
TP (mg/m <sup>3</sup> )	21.64 (1.41)	34.19 (3.56)	$P<0.01$ †
Secchi disc (m)	2.11 (0.085)	2.98 (0.12)	$P<0.0001$
TON (mg/l)	0.81 (0.1)	0.86 (0.11)	$P>0.05$

Standard error in brackets; †only 3 readings in 2000; homogeneity of variance assumptions not met for TP

Spearman's Rank correlations in the pre-zebra mussel period revealed a strong significant negative association between chlorophyll *a* and Secchi disc ( $r_s=-0.705$ ,  $df=103$ ,  $P<0.05$ ) and a weak significant negative association between TP and Secchi disc ( $r_s=-0.31$ ,  $df=94$ ,  $P<0.05$ ), whereas there was a significant, albeit weak, positive association between TP and chlorophyll *a* ( $r_s=0.33$ ,  $df=96$ ,  $P<0.05$ ). Post establishment, there was a modest significant negative association between chlorophyll *a* and Secchi disc ( $r_s=-0.55$ ,  $df=34$ ,  $P<0.05$ ), however no other significant relationships were evident between TP, chlorophyll *a* and Secchi disc. No strong or relevant associations were evident for TON in either period when correlated with the other factors (Fowler et al. 1998).

## Discussion

Overall, the zebra mussel population appears to be still expanding in Lough Sheelin as evidenced from the 2005 to 2006 data, although some local isolated declines on particular substrates were evident (e.g. at site G9 on the soft substrate). This is common in the years after colonisation although most evident during the initial exponential population growth phase (e.g. Burlakova et al. 2006; Lucy et al. 2005). Indeed, Burlakova et al. (2006) suggest that it can take from 7 to 12 years after the first introduction for a mussel population to reach maximum density in a water-body.

The biomass and density of zebra mussels on the principal substrates are within the range recorded in the literature for water bodies of similar characteristics (e.g. Burlakova et al. 2006; Stańczykowska and Lewandowski 1993). The latter is somewhat confirmed by the model developed by Ramcharan et al. (1992), where a density category <3000 mussels per m<sup>2</sup> was predicted. Although the model does not take into account substrate type, the predicted densities for 2005 and 2006 are close to the actual average density recorded on the stony substrates, but substantially higher than on the soft substrate.

The overall increase in biomass can be predominantly attributed to observed increases on the stony substrate areas of the lake in spring 2006. This was reflected in the length frequency distributions as a shift from the smaller to larger size mussels on the stony substrate (Figures 2A and 2B). On soft substrates, biomass increased at sites of 0-3m depths but a decrease was evident in the 3-5m depth range. Soft substrate sites located in the vicinity of areas of submerged vegetation growth or close to areas of stony substrate, generally had a higher abundance and biomass of zebra mussels present. Annual plant die-off from year to year most likely facilitates this, as zebra mussels fall onto the soft substrate bottom, attached to dead plant matter, potentially creating a nucleus for future colonisation by other mussels (Ramcharan et al. 1992; Burlakova et al. 2006). Furthermore, the close proximity of vegetation colonised by zebra mussels may act as sources of higher local levels of settling juvenile stages, thus enhancing associated abundance and biomass (Coakley et al. 2002). Although no zebra mussels were recorded at the deepest site in the lake (11.2m depth) in spring of either year, a subsequent diving survey in autumn 2006 confirmed that zebra mussels were present at this site, albeit at a very low percentage cover (1-3%).

The higher number of individual mussels and weight of mussels per gram of plant material found on the Characeae and *Elodea* spp. reflect the relative stability of these submerged taxa. Both are perennial and offer reasonably firm sized substrate for the attachment of juvenile mussels (Lewandowski 1982). Observed decreases in average values recorded in 2006 (see Table 1) are most likely due to the fresh plant growth evident in the 2006 samples. Although not examined, it is reasonable to assume the increase in water transparency post-establishment has resulted in the expansion of

submerged vegetation thus providing additional habitat for zebra mussels to colonise. Previous research in the lake has shown that the Characeae beds have retracted in years of low water transparency and a reduction in water quality, only to become eventually re-established after conditions improve (Champ 1993). As regards the emergent reed bed taxa examined, *Schoenoplectus lacustris* was colonised to a greater extent than *Phragmites australis*. These reeds were not as heavily colonised as those reported by Lucy (2005) and Sullivan et al. (2002) in previous Irish lake studies. However it is mentioned that the reed beds were not found to be as important as a substrate for settlement after the initial first years of population expansion (Lucy 2005).

Native unionid mussels, *Anodonta* spp., while subsequently extirpated, have proved to be a significant substrate for settlement of zebra mussels in other Irish lakes (Maguire et al. 2003; Lucy et al 2005). Although few specimens were encountered in Lough Sheelin, all had zebra mussels attached. No live specimens were recorded in the 2006 samples. However it is important to note that a number of predominantly dead *Anodonta* spp. with attached zebra mussels, have been observed retrieved in gill nets during fish population survey work in the corresponding time period. Additionally, a single live *Anodonta* mussel was found during a mid-lake scuba dive in autumn 2006 (personal observations), suggesting native mussels may still persist in the lake in the deeper soft substrate areas. As the *Anodonta* lifecycle involves a parasitic glochidial larval stage which can be associated with *Salmo trutta* Linn. (Bauer and Wächtler 2001 and references therein), the migration of these fish from the surrounding feeder rivers into the lake, may help to ensure the presence of an *Anodonta* population, albeit at a substantially lower level than the pre-invasion period.

Apart from soft substrate samples in spring 2005, length frequency distributions showed no clear cohorts to help define age classes. This is due to a number of factors, most notably the occurrence of multiple spawning events resulting in a broad range of time over which mussel settlement can potentially occur in one reproductive season. Furthermore, zebra mussel growth may not be uniform due to local temperature and food differences as well as the slowing of growth with age (reviewed in Karatayev et al. 2006).

The time (22.4 days and 13.1 days in spring 2005 and 2006 respectively) required by the

mussels (on stony and soft substrates) to filter total lake volume, may represent an underestimate of the actual time of the whole resident population, as zebra mussels on plant substrates were not included in this survey. It must be noted that the influence of seasonal spring water temperatures during sampling (7-10°C) on actual filtration rates, were not incorporated into the estimates, as there is no corresponding filtering rate provided in the literature (Karatayev et al. 1997). The pre-invasion residence time of water in the lake is believed to have been approximately six months (John et al. 1982). Therefore, these filtering estimates serve to illustrate the substantial effect the zebra mussel population may have on suppressing pelagic primary production in Lough Sheelin.

As documented with previous zebra mussel invasions (e.g. Fahnenstiel et al. 1995; Caraco et al. 1997), a significant reduction in chlorophyll *a* (a proxy measure of phytoplankton production) and increase in water transparency, concomitant with the zebra mussel invasion, is evident. However, TP levels have remained high since the mid 1990s (Kerins et al. 2007) and may even be slightly higher than the pre-invasion period. The continuing level of high phosphorus loading to the lake appears to be responsible for this (Kerins et al. 2007). This is unlike some other studies reported in the literature, in which a reduction in TP was observed post-establishment (e.g. Higgins et al. 2008; Maguire et al. 2003). TP is known to be closely related to primary production and thus chlorophyll *a* in lakes (Champ 1998). This relationship has been demonstrated previously in Lough Sheelin (Champ 1993). The positive correlation between TP and chlorophyll *a*, although statistically a relatively weak direct association, is not evident in the post establishment period. If TP continues to remain high, it may become further decoupled from chlorophyll as predicted by Kerins et al. (2007). Interestingly, a previous reduction in TP in the early 1990s corresponded with an improvement in water quality (as measured by chlorophyll *a* and Secchi disc) in the lake (Champ 1993; Champ 1998).

The three primary measures of water quality indicators in Lough Sheelin (TP, chlorophyll *a* and Secchi disc), place the lake into two separate trophic categories, eutrophic as regards TP, and mesotrophic for the latter two parameters, as assessed by current practices (OECD 1982; Toner et al. 2005). This has implications with regard to interpreting and meeting *good*

*ecological status* under the *Biological Quality Elements* as required by European Union Water Framework Directive (2000). Although the Directive makes no explicit mention of invasive species, their introduction is given as an example of a biological pressure and impact in the instructive Guidance document (2003) of the Common Implementation Strategy for the Water Framework Directive (2000/60/EC). The consideration of the influence of invasive species in general on aquatic systems has yet to be adequately resolved, although new classification systems to account for this have recently been proposed (Olenin et al. 2007).

The long-term dynamics of the resident zebra mussel population are currently unclear. Stańczykowska and Lewandowski (1993) outline 3 potential scenarios based on long-term observations of density in zebra mussel infested Polish lakes: (1) density will remain relatively stable - (large, deep and mesotrophic or moderate eutrophy lakes); (2) density will be relatively unstable with yearly fluctuations and eventual stability at generally lower densities due to increased eutrophy; and (3) rapid population crash resulting in increased eutrophication. It is worth noting that if phosphorus inputs to the lake remain high, any sudden decline in the zebra mussel population may result in the re-occurrence of significant eutrophication problems; the mussels will be unable to partake in their apparent current role to suppress primary production and act as a sizeable phosphorus sink. Additionally, this paper highlights the common misconception that zebra mussel introductions lead to overall improvements in water quality. In the case of Lough Sheelin, only an apparent improvement was evident, as TP concentrations confirm the lake to be still in a eutrophic state.

#### Acknowledgements

Special thanks to Dr Martin O'Grady of the Central Fisheries Board for his advice and for providing a basis for the project. For field support, thanks to Eamon Cusack and the staff of the Shannon Regional Fisheries Board based at Lough Sheelin, including Sean Gurhy and Martin Moffatt; members of the UCD Subaqua Club and John Kavanagh. Thanks to Dr Tasman Crowe, UCD, for some statistical advice. This research was supported by the Central and Shannon Regional Fisheries Boards, Lough Sheelin Trout

Protection Association and the Irish Research Council for Science, Engineering and Technology. Financial assistance to present this work at the 15th ICAIS conference was provided by the National Parks and Wildlife Service and the Shannon Regional Fisheries Board.

## References

- Astane J, Gosling E, Wilson J, Powell E (2005) Genetic variability and phylogeography of the invasive zebra mussel, *Dreissena polymorpha* (Pallas). *Molecular Ecology* 14(6): 1655–1666
- Bauer G, Wächtler K (eds) (2001) *Ecology and Evolution of the Freshwater Mussels Unionoida*. Springer-Verlag, Berlin, pp 424
- Burla H, Ribi G (1998) Density variation of the zebra mussel *Dreissena polymorpha* in Lake Zurich, from 1976 to 1988. *Aquatic Sciences* 60: 145–156
- Burlakova LE, Karatayev A, Padilla DK (2006) Changes in the distribution and abundance of *Dreissena polymorpha* within lakes through time. *Hydrobiologia* 571: 133–146
- Caraco NF, Cole JJ, Raymond PA, Strayer DL, Pace ML, Findlay SEG, Fischer DT (1997) Zebra mussel invasion in a large, turbid river: Phytoplankton response to increased grazing. *Ecology* 78: 588–602
- Champ WST (1977) Trophic status of fishery lakes. In: Downey WK, Ni Uid G (eds) *Lake pollution prevention by eutrophication control*. Proceedings of a seminar, Killarney 10–11 May. National Science Council, Dublin, Stationary Office, pp 65–78
- Champ WST (1993) Lough Sheelin - A 'Success' Story. In: Mollan C (ed) *Water of Life*. The proceedings of a conference on the inland waters of Ireland. Royal Dublin Society 7–9 October 1992, pp 154–162
- Champ WST (1998) Phosphorous/chlorophyll relationships in selected Irish lakes: ecological consequences and suggested criteria for ecosystem management. In: Wilson JG (ed) *Eutrophication in Irish waters*. Royal Irish Academy, pp 91–105
- Coakley JP, Rasul N, Ioannou SE, Brown GR (2002) Soft sediment as a constraint on the spread of the zebra mussel in western Lake Erie: processes and impacts. *Aquatic Ecosystem Health and Management* 5(3): 329–343
- Fahnenstiel GL, Lang GA, Nalepa TF and Johengen TH (1995) Effects of zebra mussel (*Dreissena polymorpha*) colonization on water quality parameters in Saginaw Bay, Lake Huron. *Journal of Great Lakes Research* 21: 435–448
- Fowler J, Cohen L, Jarvis P (1998) *Practical statistics for field biology*, 2nd Edition. Wiley and Sons, Chichester, England
- Gargan PG, O'Grady MF (1992) Feeding relationships of trout *Salmo trutta* L., perch *Perca fluviatilis* (L.) and roach *Rutilus rutilus* (L.) in Lough Sheelin, Ireland. *Irish Fisheries Investigations* 35A: 1–22
- Guidance document (2003) *Common implementation strategy for the Water Framework Directive (2000/60/EC). Transitional and Coastal Waters – Typology, Reference Conditions and Classification Systems*. Produced by WG 2.4. – COAST. Luxemburg, 107 pp.
- Higgins TM, Grennan JM, McCarthy TK (2008) Effects of recent zebra mussel invasion on water chemistry and phytoplankton production in a small Irish lake. *Aquatic Invasions* 3 (1): 14–20
- Hunter RD, Simons KA (2004) Dreissenids in Lake St. Clair in 2001: Evidence for population regulation. *Journal of Great Lakes Research* 30(4): 528–537
- Irvine K, Allott N, Caroni R, Eyto E de, Free G, White J (2001) Ecological assessment of Irish lakes. The development of a new methodology suited to the needs of the EU Directive on surface waters. Environmental Protection Agency, Wexford, Ireland
- John DM, Champ WST, Moore JA (1982) The changing status of Characeae in four marl lakes in the Irish midlands. *Journal of Life Sciences of the Royal Dublin Society* 4: 47–71
- Karatayev AY, Burlakova LE, Padilla DK (1997) The effects of *Dreissena polymorpha* (Pallas) invasion on aquatic communities in Eastern Europe. *Journal of Shellfish Research* 16: 187–203
- Karatayev AY, Burlakova LE, Padilla DK (2002) Impacts of zebra mussels on aquatic communities and their role as ecosystem engineers. In: Leppäkoski E, Gollasch S, Olenin S (eds) *Invasive aquatic species of Europe: Distribution, impacts and management*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 433–446
- Karatayev AY, Burlakova LE, Padilla DK (2006) Growth rate and longevity of *Dreissena polymorpha* (Pallas): a review and recommendations for future study. *Journal of Shellfish Research* 25(1): 23–32
- Karatayev AY, Lyakhovich VP, Reznichenko OG (1990) Quantitative study of the density, biomass and abundance dynamic. In: Shkhorbatov GL, Starobogatov YI (eds) *Methods for study of bivalvian molluscs*. Trudy Zoologicheskogo Instituta 219: 172–178
- Kerins C, Monahan K, Champ T (2007) Lough Sheelin and its catchment: water quality status and nutrient loadings 1998–2005. Shannon Regional Fisheries Board, Limerick, Ireland
- Lancioni T, Gaino E (2006) The invasive zebra mussel *Dreissena polymorpha* in Lake Trasimeno (Central Italy): Distribution and reproduction. *Italian Journal of Zoology* 73 (4): 335–346
- Lewandowski K (1982) The role of early developmental stages in the dynamics of *Dreissena polymorpha* (Pallas) populations in lakes. II. Settling of larvae and the dynamics of numbers of settled individuals. *Ekologia Polska* 30: 223–286
- Lovell SJ, Stone SF, Fernandez L (2006) The economic impacts of aquatic invasive species: A review of the literature. *Agricultural and Resource Economics Review* 35(1): 195–208
- Lowe RL, Pillsbury RW (1995) Shifts in benthic algal community structure and function following the appearance of zebra mussels (*Dreissena polymorpha*) in Saginaw Bay, Lake Huron. *Journal of Great Lakes Research* 21: 558–566
- Lucy F (2005) *The dynamics of zebra mussel (Dreissena polymorpha) populations in Lough Key, Co. Roscommon, 1998–2003*. PhD dissertation, Institute of Technology, Sligo
- Lucy F, Sullivan M, Minchin D (2005) Nutrient levels and the zebra mussel population in Lough Key. Environmental Research, Technological, Development and Innovation Report Series No. 34. Environmental Protection Agency, Wexford, Ireland
- Lucy F (2006) Early life stages of *Dreissena polymorpha* (zebra mussel): the importance of long-term datasets in invasion ecology. *Aquatic Invasions* 1: 171–182

## Impact of zebra mussel on Lough Sheelin

- Mackie GL (1993) Biology of the zebra mussel (*Dreissena polymorpha*) and observations of mussel colonization on unionid bivalves in lake St. Clair of the Great Lakes. In: Nalepa TF, Schloesser DW (eds) Zebra Mussels: Biology, Impacts and Control. Lewis Publishers, Boca Raton, Florida, pp 153-165
- Maguire CM (2002) The zebra mussel *Dreissena polymorpha* in the Erne system: Invasion, population dynamics and early ecological impacts. PhD thesis. The Queens University of Belfast
- Maguire CM, Roberts D, Rosell RS (2003) The ecological impact of a zebra mussel invasion in a large Irish lake, Lough Erne: a typical European experience? Aquatic Invaders 14: 2-8
- Marsden JE (1992) Standard protocols for monitoring and sampling zebra mussels. Illinois Natural History Survey Biological Notes 138, pp 40
- Marsden JE, Lansky DM (2000) Substrate selection by settling zebra mussels, *Dreissena polymorpha*, relative to material, texture, orientation, and sunlight. Canadian Journal of Zoology 78: 787-793
- Minchin D (2003) The Zebra Mussel *Dreissena polymorpha* (Pallas) extends its range westwards in Ireland. Bulletin of the Irish Biogeographical Society 27: 176-182
- Minchin D, Lucy F, Sullivan M (2002a) Zebra mussel: impacts and spread. In: Leppäkoski E, Gollasch S, Olenin S (eds) Invasive aquatic species of Europe: distribution, impact and management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 135-146
- Minchin D, Lucy F, Sullivan M (2002b) Monitoring of zebra mussels in the Shannon-Boyle navigation, other navigable regions and principal Irish lakes, 2001 and 2002. Marine Institute, Dublin, Marine Environment and Health Series No. 5
- Minchin D, Lucy F, Sullivan M (2005) Ireland: a new frontier for the zebra mussel *Dreissena polymorpha* (Pallas). Oceanological and Hydrobiological Studies 34: 19-30
- Minchin D and Moriarty C (1998a) Zebra mussels in Ireland. Fisheries Leaflet 177, Marine Institute, Dublin, Ireland
- Minchin D and Moriarty C (1998b) Distribution of the zebra mussel *Dreissena polymorpha* (Pallas) in Ireland 1997. Irish Naturalists' Journal 26: 38-42
- Munawar M, Munawar IF, Mandrak NE, Fitzpatrick M, Dermott R, Leach J (2005) An overview of the impact of non-indigenous species on the food web integrity of North American Great Lakes: Lake Erie example. Aquatic Ecosystem Health and Management 8: 375-395
- O.E.C.D. (1982) Eutrophication of waters: monitoring, assessment and control. Report of the O.E.C.D. Cooperative Programme on Eutrophication. Prepared by Vollenweider RA, Kerekes J. Organisation for Economic Cooperation and Development, Paris, pp 164
- Olenin S, Minchin D, Daunys D (2007) Assessment of biopollution in aquatic ecosystems. Marine Pollution Bulletin, Special Issue 55(7-9): 379-394
- Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs associated with alien-invasive species in the United States. Ecological Economics 52: 273-288
- Pollux B, Minchin D, Van der Velde G, Van Alen T, Moon-Van Der Staay SY, Hackstein J (2003) Zebra mussels (*Dreissena polymorpha*) in Ireland, AFLP-fingerprinting and boat traffic both indicate an origin from Britain. Freshwater Biology 48: 1127-1139
- Ramcharan CW, Padilla DK, Dodson SI (1992) Model to predict potential occurrence and density of the zebra mussel, *Dreissena polymorpha*. Canadian Journal of Fisheries and Aquatic Sciences 49(12): 2611-2620
- Reeders HH, bij De Vaate AB, Noordhuis R (1993) Potential of the zebra mussel *Dreissena polymorpha* for water quality management. In: Nalepa TF, Schloesser DW (eds) Zebra Mussels: Biology, Impacts and Control. Lewis Publishers, Boca Raton, Florida, pp 439-452
- Skubinna JP, Coon TG, Batterson TR (1995) Increased abundance and depth of submersed macrophytes in response to decreased turbidity in Saginaw Bay, Lake Huron. Journal of Great Lakes Research 21: 476-488
- Stańczykowska A, Lewandowski K (1993) Thirty years of studies of *Dreissena polymorpha* in Mazurian Lakes of northeastern Poland. In: Nalepa TF, Schloesser DW (eds) Zebra Mussels: Biology, Impacts and Control. Lewis Publishers, Boca Raton, Florida, pp 3-33
- StatSoft Inc (2008) STATISTICA (data analysis software system), version 8.0. <http://www.statsoft.com>
- Strayer DL, Malcom HM (2006) Long-term demography of a zebra mussel (*Dreissena polymorpha*) population. Freshwater Biology 51: 117-130
- Sullivan M, Lucy F, Minchin D (2002) The association between zebra mussels and aquatic plants in the Shannon River system, Ireland. Aquatic Invaders 13(4): 6-9
- Toner P, Bowman J, Clabby K, Lucey J, McGarrigle M, Concannon C, Clenaghan C, Cunningham P, Delaney J, O'Boyle S, MacCárthaigh M, Craig M, Quinn R (2005) Water Quality in Ireland 2001-2003. Environmental Protection Agency, Wexford, Ireland
- Underwood AJ (1997) Experiments in ecology: their logical design and interpretation using analysis of variance. Cambridge University Press, Cambridge, pp 504
- Vanderploeg HA, Nalepa TF, Jude DJ, Mills EL, Holeck KT, Liebig JR, Grigorovich IA, Ojaveer H (2002) Dispersal and emerging ecological impacts of Ponto-Caspian species in the Laurentian Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences 59: 1209-1228
- Water Framework Directive (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy. Official Journal of the European Communities 43: 1-72
- Wentworth CK (1922) A scale of grade and class terms for clastic sediments. Journal of Geology 30: 377-392
- Wong WH, Levinton JS, Twining BS, Fisher N (2003) Assimilation of micro- and mesozooplankton by zebra mussels: a demonstration of the food web link between zooplankton and benthic suspension feeders. Limnology and Oceanography 48(1): 308-312
- Zhu B, Fitzgerald DG, Mayer CM, Rudstam LG, Mills EL (2006) Alteration of ecosystem function by zebra mussels in Oneida Lake, NY: Impacts on submerged macrophytes. Ecosystems 9: 1017-1028